

Voronoi L-decomposition of PSD_n and the hypermetric correlation cone

Michel Deza

Ecole Normale Supérieure, Paris, France

and Viatcheslav Grishukhin

CEMI RAN, Moscow, Russia

Abstract

We describe connections of Voronoi's results with hypermetrics.

Denote by PSD_n and $HYCO_n$ the cones in $\mathbf{R}^{\binom{n+1}{2}}$ consisting of real symmetric matrices $(a_{ij})_{i,j=1}^n$ such that

$$PSD_n = \{a \in \mathbf{R}^{\binom{n+1}{2}} : \sum_{1 \leq i, j \leq n} a_{ij} x_i x_j \geq 0 \text{ for all integral } x_i\},$$

$$HYCO_n = \{a \in \mathbf{R}^{\binom{n+1}{2}} : \sum_{1 \leq i, j \leq n} a_{ij} x_i x_j - \sum_{1 \leq i \leq n} a_{ii} x_i \geq 0 \text{ for all integral } x_i\}.$$

The first cone PSD_n is non-polyhedral cone of all positive semidefinite matrices. Note that the inequalities defining PSD_n are satisfied also by all rational x , and, by continuity, by all real x . The second cone $HYCO_n$ (called the *hypermetric correlation cone*) is polyhedral. There is the third cone COR_n tightly related to these cones and called the *correlation cone*. This cone is defined not by inequalities but by its extreme rays which are all $2^n - 1$ binary matrices $(a_i(S)a_j(S))_{i,j=1}^n$ for all nonzero indicator vectors $a(S) \in \{0, 1\}^n$, $S \subseteq \{1, 2, \dots, n\}$. Hence

$$COR_n = \{a \in \mathbf{R}^{\binom{n+1}{2}} : a_{ij} = \sum_{S \subseteq \{1, \dots, n\}} x_S a_i(S) a_j(S) \text{ for all nonnegative } x_S\}.$$

COR_n and $HYCO_n$ are very interesting because images of them under some linear mapping of $\mathbf{R}^{\binom{n+1}{2}}$ correspond to semimetrics on $n+1$ points. The semimetrics of the map of COR_n are metric subspaces of some l_1^m (i.e. of \mathbf{R}^m with the metric $d_{pq} = \|p - q\|_{l_1}$). The semimetrics of the map of $HYCO_n$ are *hypermetrics* $(d_{ij})_{i,j=1}^{n+1}$, i.e. they satisfy the following *hypermetric inequalities*:

$$\sum_{1 \leq i < j \leq n+1} b_i b_j d_{ij} \leq 0 \text{ for all } b \in \mathbf{Z}^{n+1} \text{ with } \sum_{1 \leq i \leq n+1} b_i = 1.$$

See book [DL97] for detailed study of those semimetrics; see, especially, Chapters 13–17 there.

The purpose of this note is to present the cone $HYCO_n$ (especially its polyhedrality) briefly and only in the context of the remarkable decomposition of PSD_n into polyhedral L-domains and the finiteness of the number of types of L-polytopes given in the famous Deuxième mémoire of Voronoi published in 1908 and (the second part) in 1909 by Crelle Journal.

First, we mention that $HYCO_n = COR_n$ iff $n \leq 5$ (see [De60] for $n = 4$ and [Ba71] for $n = 5$). The set of all facets of $HYCO_6$ is given in [Ba95]. The number $O(n)$ of orbits of facets under the permutation group S_n , and the number of all facets of $HYCO_n$ are as follows

n	2	3	4	5	6
$O(n)$	1	1	2	4	14
$N(n)$	3	12	40	210	3773

Now we will to connect the polyhedrality of $HYCO_n$ with the following two theorems of Voronoi; their proof is, actually, the main content of Deuxième mémoire [Vor08], [Vor09].

Theorem A (Voronoi) *The action of the group $GL(n, \mathbf{Z})$ induces a partition of the cone PSD_n into disjoint relatively open convex subcones, called the L-type domains, of dimension $1, 2, \dots, \binom{n+1}{2}$, and having the following properties:*

(i) *On each of these subcones the affine structure of the L-decomposition of corresponding lattices is constant.*

(ii) *Subcones of dimension $\binom{n+1}{2}$ correspond to general lattices, i.e. having simplicial L-decompositions. These L-type domain are polyhedral.*

(iii) *A subcone of dimension less than $\binom{n+1}{2}$ is a relatively open face of two or more L-type domains. If such a cone makes contact with the boundary of an L-type domain, then it is necessarily a face of that domain. The lattice corresponding to a quadratic form on such a face is special, i.e. it has among its Delaunay polytopes some that are not simplices.*

Theorem B (Voronoi) *The numbers of distinct L-types (i.e. the numbers of stars of L-polytopes) of lattices in \mathbf{R}^n is finite for each dimension n .*

Our main theorem is that $HYCO_n$ is polyhedral. The first proof ([DGL93]) used Theorem B. Since Voronoi’s proof of Theorem B was very involved, we gave also in [DGL93] much shorter (2 pages) proof of Theorem B. Another proof of the polyhedrality of $HYCO_n$ follows from the Theorem A of Voronoi. In fact, let $v_1, \dots, v_n \in \mathbf{R}^k$, where $1 \leq k \leq n$. We set $L =: \mathbf{Z}(v_1, \dots, v_n)$ and define $p \in PSD_n$ by setting $p_{ij} = v_i^T v_j$ for $1 \leq i, j \leq n$. Then

(*) $p \in HYCO_n$ if and only if L is a lattice and v_1, \dots, v_n are all vertices of the same Delaunay polytope in the star of L at the origin.

Moreover, if $p \in HYCO_n$, then the whole L-type domain containing p is entirely contained in $HYCO_n$. Hence $HYCO_n$ is a union of L-type domains. In fact, this union is *finite*. Indeed, $HYCO_n$ contains only finitely many L-type domains whose associated lattices have the same L-type (by (*)) and since, in a given lattice L , there are only finitely many ways of choosing a set of n vectors v_1, \dots, v_n that all are vertices of the same Delaunay polytope in the star of L . Finally, the number of distinct L-types of lattices of

given dimension is finite. Therefore, $HYCO_n$ is a finite union of L-type domains. But, a fundamental property is that each L-type domain is a polyhedral cone. Hence $HYCO_n$ is a polyhedral cone.

The best known upper bound on the number of facets of $HYCO_n$ (using a result of [Lo94]) is $(1 + 2^{n+1}n!/\binom{2n}{n})^n$. So, the problem of testing membership in $HYCO_n$ is in *co-NP*, but it is not known whether this problem is *NP*-hard. In [AG93], more precise complexity results are given for integer-valued members of $HYCO_n$.

L-polytopes were introduced in the Deuxième mémoire of Voronoi; they also called Delaunay polytopes, since Delaunay and his school worked a lot on them, see the introduction of [NRDS92] for a historical account; see also [Er92], [RB79]. The link between hypermetrics and L-polytopes was established in [As82]. L-polytopes versus perfect lattices were considered in [Gr93], versus equiangular lines, in [DG95a], versus covering radius of a lattice, in [DG96]. Low-dimensional L-polytopes were considered in [ER87] and [DG95b]. Subclasses of members of $HYCO_n$ (coming from integer-valued graphic hypermetric) were considered in [TD87] and [DG93].

Besides the polyhedrality of $HYCO_n$, the main notion introduced in [DGL92], [DGL93], [DGL95] is the *rank* of an L-polytope, and we hope that it will be useful in the geometry of quadratic forms. In short, the rank of an L-polytope is the number of its degrees of freedom that it has when one deforms it in such a way that the deformed polytope remains an L-polytope. More precisely, the rank $rk(P)$ of a Delaunay polytope P is the dimension (in a topological sense) of the set of all affine bijections T of \mathbf{R}^k (if P is k -dimensional) up to translations and orthogonal transformations, such that $T(P)$ is again an L-polytope, see [La96]. The main results about the rank given in [DGL92], [DGL93], [DGL95] are:

- 1) $rk(P)$ is equal to the rank of any generating subset of the set $V(P)$ of vertices of P ;
- 2) $rk(P_1 \times P_2) = rk(P_1) + rk(P_2)$;
- 3) Suppose that an n -dimensional L-polytope P is *basic*, i.e. $V(P)$ contains a basis of the lattice generated by $V(P)$; all known L-polytopes are basic. Then

$$\binom{n+2}{2} - |V(P)| \leq rk(P) \leq \binom{n+1}{2}$$

and, moreover, $\binom{n+1}{2} - \frac{|V(P)|}{2} + 1 \leq rk(P)$ if P is centrally symmetric. For example, the simplex α_n realizes both the inequalities in the first bound; the cross-polytope β_n realizes the equality in the second bound on the rank of a symmetric Delaunay polytope.

- 4) The *repartitioning* polytopes, introduced in the Deuxième mémoire of Voronoi, but named and studied in [RB79], are exactly L-polytopes of *corank* 1, i.e. of rank $\binom{n}{2} - 1$, see [AG93].

We call L-polytopes of rank 1 *extreme*. An extreme L-polytope is rigid in the sense that a homotety is the only affine transformation of it which is still an L-polytope. In terms of the correspondence between L-polytopes and hypermetrics, the above two cases of Delaunay polytopes of maximal and minimal rank correspond to facets and extreme rays of $HYCO_n$. Examples of extreme L-polytopes, given in [DGL92], [DGL93], [DGL95], are:

the segment α_1 , the Schläfli $2_{21} \in \mathbf{R}^6$ and the Gosset $3_{21} \in \mathbf{R}^7$ polytopes (constructed from sections of the root lattice E_8); two centrally symmetric polytopes in \mathbf{R}^{16} and 3 polytopes in \mathbf{R}^{15} (from sections of the Barnes-Wall lattice Λ_{16}); a centrally symmetric L-polytope in \mathbf{R}^{23} with 552 vertices and an asymmetric L-polytope in \mathbf{R}^{22} with 275 vertices (from the Leech lattice Λ_{24}).

It will be interesting to construct an infinite sequence of extreme L-polytopes.

For any n , $HYCO_n$ and COR_n have $2^n - 1$ common extreme rays $(a_i(S)a_j(S))_{i,j=1}^n$, which are preimages of the cut metrics. These are only extreme rays for $n \leq 5$. We know ([DGL92], [DGL93]) additionally 26 orbits of the group S_6 of extreme rays of $HYCO_6$ distinct from cut rays. We expect that there are no other extreme rays. In this case, the segment α_1 and the Schläfli polytope 2_{21} are only Delaunay polytopes of rank 1 having dimension at most 6.

References

- [As82] P.Assouad, *Sous espaces de L^1 et inégalité hypermétriques*, Comptes Rendus de l'Académie de Sci. Paris **294**(A) (1982) 439–442.
- [AG93] D.Avis and V.P.Grishukhin, *A bound on k -gonality of facets of the hypermetric cone and related complexity problems*, Comput. Geometry: Theory and Appl. **2** (1993) 241–254.
- [Ba71] E.P. Baranovskii, *Simplexes of L -subdivisions of Euclidean spaces*, Math. Notes **10** (1971) 827–834 (Translated from: Math. Zametki **10** (1971) 659–671).
- [Ba95] E.P.Baranovskii, In the Second International Conference "Algebraic, Probabilistic, Geometrical, Combinatorial and Functional Methods in the Theory of Numbers", September 1995, Voronezh, Russia.
- [De60] M.E.Tylkin (=M.Deza), *On Hamming geometry of unitary cubes* (in Russian) Doklady Akad. Nauk SSSR **134** (1960) 1037–1040 (Translated in: Cybernetics and Control Theory **134** (1961) 940–943).
- [DG93] M.Deza, V.P.Grishukhin, *Hypermetric graphs*, The Quart. J. of Math. Oxford (2) **44** (1993) 399–433.
- [DG95a] M.Deza and V.P.Grishukhin, *L -polytopes and equiangular lines*, Discrete Appl. Math. **56** (1995) 181–214.
- [DG95b] M.Deza and V.P.Grishukhin, *Delaunay polytopes of cut lattices*, Linear Algebra and its Appl. **226** (1995) 667–686.
- [DG96] M.Deza and V.P.Grishukhin, *Bounds on the covering radius of a lattice*, Mathematica **43** (1996) 159–164.

- [DGL92] M.Deza, V.P.Grishukhin and M.Laurent, *Extreme hypermetrics and L-polytopes*, in: G.Halász et al., editors, *Sets, Graphs and Numbers, Budapest, (Hungary), 1991*, vol.60 of Colloquia Math. Soc. János Bolyai (1992) 157–209.
- [DGL93] M.Deza, V.P.Grishukhin and M.Laurent, *Hypermetric cone is polyhedral*, *Combinatorica* **13** (1993) 397–411.
- [DGL95] M.Deza, V.P.Grishukhin and M.Laurent, *Hypermetrics in geometry of numbers*, in: W.Cook, L.Lovasz and P.D.Seymour, editors, *Combinatorial Optimization*, vol. 20 of DIMACS Series in Discrete Math. and Theor. Comput. Sci. (1995) 1–109.
- [DL97] M.Deza and M.Laurent, *Geometry of cuts and metrics*, Springer-Verlag, Berlin, 1997.
- [Er92] R.M.Erdahl, *A cone of inhomogeneous second-order polynomials*, *Discrete and Comput. Geometry* **8** (1992) 387–416.
- [ER87] R.M.Erdahl and S.S.Ryshkov, *The empty sphere*, *Canad. J. Math.* **39** (1987) 794–824.
- [Gr93] V.P.Grishukhin *L-polytopes, even unimodular lattices and perfect lattices*, Rapport LIENS-93-1, Ecole Normale Supérieure, Paris, 1993.
- [La96] M.Laurent, *Delaunay transformations of a Delaunay polytope*, *J. of Algebraic Combinatorics* **5** (1996) 37–46.
- [Lo94] L.Lovász, Personal communication, 1994.
- [NRDS92] S.P.Novikov, S.S.Ryshkov, N.P.Dolbilin and M.A.Shtan’ko, editors, *Discrete Geometry and Topology*, Proceedings of the Steklov Institute of Math. vol. 196, 1992.
- [RB79] S.S.Ryshkov and E.P.Baranovskii, *C-types of n-dimensional lattices and 5-dimensional primitive parallelohedra (with applications to the theory of covering)*, *Trudy Math. Steklov’s Institute* **137** (1976) = Proceedings of the Steklov Institute of Mathematics (1978, No4).
- [RB79] S.S.Ryshkov and E.P.Baranovskii, *Classical methods in the theory of lattice packings*, *Russian Math. Surveys* **34**(4) (1979) 1–68.
- [TD87] P.Terwiliger and M.Deza, *Classification of finite connected hypermetric spaces*, *Graphs and Combinatorics* **3** (1987) 293–298.
- [Vor08] G.F.Voronoi, *Nouvelles applications des paramètres continus à la théorie des forms quadratiques - Deuxième mémoire*, *J. für die Reine und Angewandte Math.* **134** (1908) 198–287.
- [Vor09] G.F.Voronoi, *Nouvelles applications des paramètres continus à la théorie des forms quadratiques - Deuxième mémoire - Seconde partie*, *J. für die Reine und Angewandte Math.* **136** (1909) 67–181.